			1
Student's Name:	D.O.B:Teacher:		Place Child'
ALLERGY TO:_			Pictur
Asthmatic Yes*	No *Higher risk for severe reaction		Here
220			
G	◆ STEP 1: TREATMENT ◆	Ci Chld	N/ - 2' - 4' 4'
Symptoms:		**(To be determined by treatment)	y physician authorizing
■ If a food a	llergen has been ingested, but no symptoms:		☐ Antihistamine
Mouth	Itching, tingling, or swelling of lips, tongue, mouth	1 1	☐ Antihistamine
■ Skin	Hives, itchy rash, swelling of the face or extremities	☐ Epinephrine 【	☐ Antihistamine
■ Gut	Nausea, abdominal cramps, vomiting, diarrhea		☐ Antihistamine
■ Throat†	Tightening of throat, hoarseness, hacking cough	☐ Epinephrine 【	☐ Antihistamine
■ Lung†	Shortness of breath, repetitive coughing, wheezing	☐ Epinephrine 【	☐ Antihistamine
■ Heart†	Weak or thready pulse, low blood pressure, fainting, pale, blueness	☐ Epinephrine 【	Antihistamine
■ Other†		☐ Epinephrine 【	Antihistamine
) Other			
If reaction DOSAGE Epinephrine: injection	tis progressing (several of the above areas affected), give: †Potentially life-threatening. The severity of symptoms can quickly continuous culture of the severity of symptoms can also continuous culture of the severity of symptoms can continuous culture of the severity culture of the severity of symptoms can continuous culture of the severity of symptoms can continuous culture of the severity of symptoms can continuous culture of the severity culture of the severi	hange.	□ Antihistamine
DOSAGE Epinephrine: injective (see reverse side for	†Potentially life-threatening. The severity of symptoms can quickly continuous cularly (circle one) EpiPen® EpiPen® Jr. Twinject@or instructions)	hange.	
DOSAGE Epinephrine: injective (see reverse side for	†Potentially life-threatening. The severity of symptoms can quickly continuous cut intramuscularly (circle one) EpiPen® EpiPen® Jr. Twinject@or instructions)	hange.	
DOSAGE Epinephrine: inject (see reverse side for Antihistamine: given	†Potentially life-threatening. The severity of symptoms can quickly continuous cularly (circle one) EpiPen® EpiPen® Jr. Twinject@or instructions) ve	hange.	
DOSAGE Epinephrine: inject (see reverse side for Antihistamine: give Other: give	†Potentially life-threatening. The severity of symptoms can quickly continuous cularly (circle one) EpiPen® EpiPen® Jr. Twinject@or instructions)	hange. © 0.3 mg Twinject@	® 0.15 mg
DOSAGE Epinephrine: inject (see reverse side for Antihistamine: give Other: give	†Potentially life-threatening. The severity of symptoms can quickly continuous cularly (circle one) EpiPen® EpiPen® Jr. Twinject® or instructions) ve	hange. 3 0.3 mg Twinject@	® 0.15 mg
DOSAGE Epinephrine: inject (see reverse side for Antihistamine: give Other: give IMPORTANT: Antihistamine: Antihistamine: give IMPORTANT: give IMPORTANT	†Potentially life-threatening. The severity of symptoms can quickly continuous cularly (circle one) EpiPen® EpiPen® Jr. Twinject@ or instructions) we	hange. 3 0.3 mg Twinject@	® 0.15 mg
DOSAGE Epinephrine: inject (see reverse side for Antihistamine: give IMPORTANT: A. 1. Call 911 (or Resonant)	†Potentially life-threatening. The severity of symptoms can quickly continuous cularly (circle one) EpiPen® EpiPen® Jr. Twinject® or instructions) ve	o replace epinephred, and additional epin	© 0.15 mg ine in anaphyla
DOSAGE Epinephrine: inject (see reverse side for Antihistamine: give IMPORTANT: And 1. Call 911 (or Rescaled).	†Potentially life-threatening. The severity of symptoms can quickly continuous cularly (circle one) EpiPen® EpiPen® Jr. Twinject® or instructions) we	o replace epinephred, and additional epin	® 0.15 mg ine in anaphyla
DOSAGE Epinephrine: inject (see reverse side for Antihistamine: give IMPORTANT: A. 1. Call 911 (or Rescalation) 2. Dr 3. Parent	†Potentially life-threatening. The severity of symptoms can quickly content intramuscularly (circle one) EpiPen® EpiPen® Jr. Twinject® or instructions) we	o replace epinephred, and additional epin	® 0.15 mg ine in anaphyla
DOSAGE Epinephrine: inject (see reverse side for Antihistamine: give Other: give IMPORTANT: Antihistamine: give 1. Call 911 (or Rescalations) 3. Parent 4. Emergency contangle (see reverse side for Antihistamine: give) 4. Emergency contangle (see reverse side for Antihistamine: give) 4. Emergency contangle (see reverse side for Antihistamine: give) 4. Emergency contangle (see reverse side for Antihistamine: give) 4. Emergency contangle (see reverse side for Antihistamine: give)	†Potentially life-threatening. The severity of symptoms can quickly content intramuscularly (circle one) EpiPen® EpiPen® Jr. Twinject® or instructions) ve	o replace epinephred, and additional epin	ine in anaphyla
DOSAGE Epinephrine: inject (see reverse side for Antihistamine: give Other: give IMPORTANT: A. 1. Call 911 (or Rescalations) 2. Dr. 3. Parent 4. Emergency contangle Name/Relationships a.	†Potentially life-threatening. The severity of symptoms can quickly content intramuscularly (circle one) EpiPen® EpiPen® Jr. Twinject® or instructions) we	to replace epinephred, and additional epin	® 0.15 mg ine in anaphyla

(Required)